

**Taxonomic studies on the genus *Grandidierella* Coutière  
(Crustacea, Amphipoda)  
II. The Malagasy species**

by Alan A. MYERS \*

**Abstract.** — A review is given of the Malagasy species of the amphipod crustacean genus *Grandidierella* Coutière. *G. spinicoxa* sp. nov. is described and figured and its relationship to *G. mahafalensis* Coutière is discussed.

**Résumé.** — Étude d'une petite collection de *Grandidierella* de Tuléar, sud-ouest de Madagascar. Il apparaît que le matériel présente de grandes affinités avec *G. mahafalensis* Coutière, mais qu'il en diffère par certains caractères importants. Il est complètement décrit et figuré sous le nom de *G. spinicoxa* sp. nov. L'espèce correspond aux spécimens brièvement décrits par BARNARD (1958) de Mohéli (Comores), ce qui montrerait sa large distribution. Les autres représentants du genre signalés de Madagascar sont mentionnés.

---

INTRODUCTION

The genus *Grandidierella* Coutière (1904) was described to accommodate a species of amphipod (*G. mahafalensis* Coutière) from the land-locked saline lake Tsimanampetsotsa in the Mahafaly region of S-W Madagascar. Coutière (1904) remarked that it would be most interesting to compare his material with marine examples from the adjoining West Coast, but although the genus has been rather widely studied in the world tropics, little work has been carried out on the Malagasy species, apart from that of MONOD (1935), RUFFO (1958) who first described *G. bonnieroides* Stephensen (as *G. Bonnierii*) from Soalara and LEDOYER (1967, 1968, 1969).

Recently I received from Pr. Th. MONOD, a small collection of a species of *Grandidierella* from Tuléar in S-W Madagascar, together with topotypic material of *G. mahafalensis*. The Tuléar material shows close affinity with *G. mahafalensis* but differs in a number of important characters. It does, however, agree with material rather briefly described by BARNARD (1958) from Mohéli in the Comoro islands, which suggests the possibility of a wide distribution for this species at least along the Western coast of Madagascar and the islands of the Mozambique channel. The dissimilarity of the isolated lake Tsimanampetsotsa material from that of the widely separate but continuous marine environments at Tuléar and Mohéli would be expected, but whether the long isolation in the lake<sup>1</sup> has resulted in full speciation or merely sub-speciation is impossible to ascertain without

\* Department of Zoology, University College, Cork, Eire.

1. According to Pr. René BATTISTINI, the age of lake Tsimanampetsotsa is at least 75,000 years.

further research on living material. Considering the constant differences between Tuléar and lake Tsimanampetsotsa material it would seem most practical to describe the fully marine material as a new species, here proposed as *G. spinicoxa* sp. nov., whilst at the same time noting the close affinity of the two species.

## THE MALAGASY SPECIES

### ***Grandidierella bonnieroides* Stephensen**

*Grandidierella bonnieroides* Stephensen, 1948 : 12-16, fig. 3 ; MYERS, 1970 : 141, fig. 1-2.  
*Grandidierella bonnieri* RUFFO, 1958 : 58, fig. 8-9 ; LEDOYER, 1967 : 137, fig. 28a ; 1968 : 53, fig. 25b  
(Non) *Grandidierella bonnieri* Stebbing, 1908 : 120-123, pl. 6.

DISCUSSION. — This species is well described and figured from Madagasear by RUFFO (1958) and LEDOYER (1967). For a discussion and revised synonymy of this species see MYERS (1970).

Recorded distribution in Madagasear : Soalara (RUFFO, 1958), Tuléar (LEDOYER, 1967).

### ***Grandidierella grossimana* Ledoyer**

*Grandidierella grossimana* Ledoyer, 1968 : 53, fig. 25a.

DISCUSSION. — *G. grossimana* differs from all other Malagasy species in the structure of the male gnathopod II in which the carpus is relatively short and broad, the posterodistal margin produced into a weakly developed tooth, and the propodus broad, approximately parallel sided with a transverse palm. Other diagnostic features include the configuration of the male gnathopod I and the structure of the antennule (for which see LEDOYER, 1968).

Recorded distribution in Madagasear : Tuléar.

### ***Grandidierella mahafalensis* Coutière**

*Grandidierella mahafalensis* Coutière, 1904 : 166-170, fig. 1-19 ; RUFFO, 1958 : 55, fig. 7-8.  
*Grandidierella megnae* MONOD, 1935 : 464-465.  
(Non) *Grandidierella megnae* (Giles), 1888 : 231-235, pl. 8, fig. 1-4.

DISCUSSION. — For a recent review and appraisal of this species see RUFFO (1958). The female gnathopod I figured by RUFFO, 1958 (fig. 7, 4-5), does not appear to be of the usual form for the female of this species, more closely resembling the developing gnathopod of a young male. Since the gnathopod is from an ovigerous female it probably represents an intersex. Normal female and male gnathopods, male eoxae I and II and uropod III




FIG. 1. — *Grandidierella mahafalensis* Coutière, Lake Tsimanampetsotsa.  
 A, ♀ Gnathopod II; B, ♀ Gnathopod I; C, Uropod III; D, ♂ Gnathopod II; E, ♂ Gnathopod I; F, ♂  
 Coxae I & II.

are figured here (fig. 1 A-F) for comparison with the corresponding appendages of *G. spinicoxa* sp. nov.

Recorded distribution in Madagascar : Lake Tsimanampetsotsa.

***Grandidierella spinicoxa* sp. nov.**

*Grandidierella mahafalensis* BARNARD, 1958 : 105, fig. 19.

*Grandidierella bonnieri* LEDOYER, 1969 : 186, fig. 3.

(Non) *Grandidierella bonnieri* Stebbing, 1908 : 120-123, pl. 6.

*Description of male.* — Length 8.0 mm ; *head* shorter than first two pereon segments combined, lateral cephalic lobes well developed, rounded, eye ovoid ; *pereon* segment I with a well developed, forwardly curving, mid-ventral, sternal tooth ; *antenna* I about equal to the body length, basi-distal ratio of peduncular articles, approximately 5:6:1, flagellum longer than peduncle, with 25-30 articles, accessory flagellum uni-articulate, about equal to one half the length of the first flagellar article of the primary flagellum ; *antenna* II subpediform, two thirds the length of antenna I, peduncular article 4, longer and broader than 5, flagellum less than one half the length of the fifth peduncular article, with about 7 articles, the basal articles scarcely constricted ; *mouthparts* not differing significantly from those of *G. mahafalensis* ; *gnathopod* I coxa shallow, the anterior distal corner produced into a rounded tooth, basis robust, the posterior margin sinuous, carpus subovoid, the posterior distal angle produced into a well developed outwardly deflected tooth, a second small blunt tooth on the distal margin, posterior to the inserted propodus, and a further, more slender tooth on the posterior margin, the posterior margin between the posterior marginal tooth and the distal tooth, irregularly serrate, propodus slender, short, broader distally, the palmar angle produced into a forwardly projecting rounded tooth, dactylus equal in length to the propodus and hence greatly overlapping palm, the posterior margin produced basally into a blunt lobe, and with a small tooth midway between lobe and tip ; *gnatopod* II coxa subtriangular, the anterior distal corner produced into a small pointed tooth, basis elongate and slender, merus with a distal row of long pectinate setae, carpus elongate, about three times as long as broad, the anterior margin bearing a 'brush' of very fine setae inserted into relatively large and conspicuous bases, propodus about two thirds the length of the carpus with a series of spines on the distal portion of the posterior margin, palm almost transverse, dactylus overlapping palm ; *pereopods* III-VII scarcely differing from those of *G. mahafalensis* ; *uropod* I peduncle equal in length to the rami, the ventral distal margin produced into a curved tooth, rami subequal, spiniferous on the anterior dorsal margin ; *uropod* II peduncle shorter than rami, the outer ramus slightly the longer, both rami spiniferous on the anterior dorsal margin ; *uropod* III peduncle very short, the posterior (inner) margin

FIG. 2. — *Grandidierella spinicoxa* sp. nov., Tuléar.

A, ♂ Head and pereon segments I & II, with a, coxae I & II, b, accessory flagellum of antenna I and c, flagellum of antenna II ; B, ♂ Gnathopod I with d, palm of propodus ; C, ♂ Gnathopod II (distal-most setae of anterior margin of carpus omitted to show bases), with e, palm of propodus ; D, ♀ Gnathopod II.



expanded into a flange, the single ramus slightly over twice the length of the peduncle, with a series of stout setae on the anterior and posterior margins and a terminal group of stout setae; *telson* similar to that of *G. mahafalensis*.

*Description of female.* — Length 7.0 mm; similar to the male, but *pereon* segment I without a sternal tooth, antennae shorter, *antenna* I equal to about one half the body length, primary flagellum with 16-20 articles; *antenna* II about two thirds the length of antenna I; *gnathopod* I coxa shallow, unproduced, basis slender, about three times as long as broad, carpus about two times as long as broad, propodus shorter than the carpus with a series of moveable spines on the posterior margin, dactylus robust, overlapping palm; *gnathopod* II coxa subtriangular, unproduced, basis slender, merus with a distal row of long pectinate setae, carpus slender with a single row of long setae on the anterior margin, propodus slightly shorter than the carpus, dactylus fitting palm.

Type locality: Tuléar, S-W Madagascar.

Material examined: 4 ♂, 3 ♀.

Holotype: M.N.H.N., n° 7229.

DISCUSSION. — *G. spinicoxa* sp. nov. is closely related to *G. mahafalensis* Coutière, but differs in the following important characters: Male gnathopod I carpus more ovoid, the posterior distal margin strongly serrate, propodus with a well developed palmar tooth, dactylus euryly lobed and toothed on the posterior margin. Male gnathopod II with a 'brush' of fine setae on the anterior margin, each seta with a large, conspicuous round point of insertion. Male coxae I and II each with a distinct tooth at its anterior distal corner. Uropod III ramus broader with more robust, stout setae.

LEDOYER (1969) noted that not all the specimens in his sample of *G. spinicoxa* (as *G. bonnieri*) had well developed eoxal teeth and suggested that this was a case of polymorphism. It seems more likely, however, that those specimens with eoxal teeth represent terminal males, those without teeth, the non-terminal males, since male amphipods continue to differentiate morphologically long after attaining sexual maturity. In Aoridae and related families, terminal males may be rare or even absent from an entire population, and this occurrence of 'neotonous' populations presents considerable difficulties to taxonomic studies.

Ecology: In the inter-tidal zone among the sea grasses *Syringodium isoetifolium*, *Halophila minor*, *Diplanthera beaudettei*, *D. uninervis*, *Zostera capensis*, *Cymodocea serrulata* and *Thalassia* spp.

Distribution: Tuléar (Madagascar), Mohéli (Comoro Islands).

#### Acknowledgments

I am indebted to Pr. Th. Monod for drawing my attention to the problem and for providing me with material upon which to base the present study.




FIG. 3. — *Grandidierella spinicoxa* sp. nov., Tuléar.

A, ♂ Uropod I; B, ♂ Uropod III & Telson; C, ♂ Uropod II; D, Maxilliped; E, Labium; F, Mandible; G, Maxilla I; H, ♀ Gnathopod I; I, Maxilla II.

## REFERENCES

- BARNARD, K. H., 1958. — Terrestrial isopods and amphipods from Madagascar. *Mém. Inst. scient. Madagascar*, sér. A, **12** : 67-111.
- COUTIÈRE, H., 1904. — Sur un type nouveau d'Amphipode, *Grandidierella mahafalensis*, provenant de Madagascar. *Bull. Soc. philomath. Paris*, sér. 9, **6** : 166-174, fig. 1-19.
- GILES, G. M., 1888. — Further notes on the Amphipoda of Indian waters. *J. Asiat. Soc. Beng.*, **57** : 220-255.
- LEDOYER, M., 1967. — Amphipodes gammariens des herbiers de phanérogames marines de la région de Tuléar (République malgache). Étude systématique et écologique. *Annls Fac. Sci. Univ. Madagascar*, **5** : 124-170.
- 1968. — Amphipodes gammariens de quelques biotopes de substrat meuble de la région de Tuléar. *Annls Fac. Sci. Univ. Madagascar*, **6** : 17-62, fig. 1-25.
- 1969. — Amphipodes gammariens du sédiment des herbiers de phanérogames marines et des dunes hydrauliques du Grand récif de Tuléar (Madagascar). Étude systématique et écologique. *Recl Trav. Stn. mar. Endoume*, fasc. h.s., **9** : 183-191.
- MONOD, Th., 1935. — Contribution à l'étude faunistique de la réserve naturelle du Manampetsa (Madagascar), B. Crustacés. *Ann. Sci. nat. Paris*, sér. 10, **18** : 449-466.
- MYERS, A. A., 1970. — Taxonomic studies on the genus *Grandidierella* Coutière (Crustacea : Amphipoda), with a description of *G. dentimera* sp. nov. *Bull. mar. Sci. Gulf Caribb.*, **20** : 135-147.
- RUFFO, S., 1958. — Amphipodes terrestres et des eaux continentales de Madagascar, des Comores et de la Réunion. *Mém. Inst. scient. Madagascar*, sér. A, **12** : 35-66.
- STEBBING, T. R. R., 1908. — The fauna of brackish ponds at Port Canning, Lower Bengal. Part 9 : A new species of Amphipoda. *Rec. Indian Mus.*, **2** (2) : 119-123.
- STEPHENSEN, K., 1948. — Amphipods from Curaçao, Bonaire, Aruba and Marguerita. *Stud. Fauna Curaçao*, **3** (11) : 1-20.

*Manuscrit déposé le 26 août 1971.*

*Bull. Mus. Hist. nat., Paris*, 3<sup>e</sup> sér., n<sup>o</sup> 64, juillet-août 1972,  
Zoologie 50 : 789-796.

*Achévé d'imprimer le 30 décembre 1972.*